

A NOTE ON A CONJECTURE OF L. J. MORDELL

BY

MICHAELA. MALCOLM

STAN-CS-70-184
NOVEMBER 1970

COMPUTER SCIENCE DEPARTMENT
School of Humanities and Sciences
STANFORD UNIVERSITY



A NOTE ON A CONJECTURE OF L. J. MORDELL

by

Michael A. Malcolm

Abstract: A computer proof is described for a previously unsolved problem concerning the inequality

$$\sum_{i=1}^n x_i / (x_{i+1} + x_{i+2}) \geq \frac{n}{2} .$$

A NOTE ON A CONJECTURE OF L. J. MORDELL

by

Michael A. Malcolm */

The function

$$S_n(\underline{x}) = \sum_{i=1}^n \frac{x_i}{x_{i+1} + x_{i+2}}, \quad (\text{subscripts mod } n) \quad (1)$$

where $x_i > 0$, $x_i + x_{i+1} > 0$, $i = 1, 2, \dots \pmod{n}$, has been studied by Shapiro, Mordell, and others (see [1-2, 6-11]). Let

$$\lambda(n) = \inf_x S_n(\underline{x}). \quad (2)$$

Then

$$h(n) \leq n/2. \quad (3)$$

H. S. Shapiro [10] suggested the verification of

$$h(n) = n/2. \quad (4)$$

For $n \leq 6$, several authors (see [6]) proved the validity of (4).

Mordell [6] conjectured that (4) is false for all $n \geq 7$, but later [7]

stated that (4) is true for $n = 7$. M. J. Lighthill (see [9]) and

A. Zulauf [11] proved that (4) is false for $n = 2k$, $k \geq 7$.

D. Z. Djokovic [2] proved that (4) is true for $n = 8$. P. H. Diananda [1]

proved that (4) is false for $n = 2k+1$, $k \geq 13$. In the same paper he

*/ This work was sponsored in part by the Office of Naval Research (NR 044-211) and by the National Science Foundation (GJ 408).

proved (i) that if (4) is false for some $n = m$, where m is odd, then (4) is false for all $n \geq m$, and (ii) that if (4) is true for some $n = m$, where m is even, then (4) is true for all $n < m$. Recently, P. Nowosad [8] proved (4) is true for $n = 10$, and thus (4) is true for all $n \leq 10$.

Therefore, the question remains open for $n = 12$ and for odd n ranging from 11 to 25. In this note I will describe a computer proof that (4) is false for $n = 25$.

The problem can be viewed as a multivariate constrained minimization of s_n . The constraints $x_i \geq 0$, $i = 1, 2, \dots, n$, can be removed by the simple change of variables

$$x_i = \theta_i^2, \quad i = 1, 2, \dots, n.$$

If the second constraint is violated, i.e., $x_i + x_{i+1} = 0$ for some i , the function (1) is undefined. Thus, the function

$$s_n(\theta) = \sum_{i=1}^n \frac{\theta_i^2}{\theta_{i+1}^2 + \theta_{i+2}^2} \quad |$$

where the indices are taken mod n , can be minimized using a standard method. For the case $n = 25$, this was done using a subroutine written by J. Alan George [3] on an IBM 360/67. The resulting values for x_i were:

$x_1^* = 5.71138$	$x_{13}^* = 2.47011$
$x_2^* = 0$	$x_{14}^* = 3.15375$
$x_3^* = 6.76097$	$x_{15}^* = 1.66622$
$x_4^* = 1.10052$	$x_{16}^* = 3.058$
$x_5 = 6.90241$	$x_{17}^* = 0.980738$
$x_6^* = 2.57379$	$x_{18}^* = 3.12582$
$x_7 = 5.91561$	$x_{19}^* = 0.400648$
$x_8^* = 3.33613$	$x_{20}^* = 3.44328$
$x_9^* = 4.60951$	$x_{21}^* = 0$
$x_{10}^* = 3.47149$	$x_{22}^* = 4.07589$
$x_{11}^* = 3.43693$	$x_{23}^* = 0$
$x_{12}^* = 3.33360$	$x_{24}^* = 4.8248$
	$x_{25}^* = 0$

To prove that $S_{25}(x^*)$ is less than 12.5 , the calculation (1) was programmed in a language [4] in which the calculations are carried out in interval arithmetic (see Moore [5]). The inaccuracies due to number conversion and **roundoff** are automatically accounted for by the language translator. Thus, assuming the translator is properly programmed, the resulting interval is guaranteed to contain the true result. The program was run on the IBM 360/67 at Stanford University, giving

$$12.49847 < S_{25}(x^*) < 12.49851 .$$

Computer time, including program compilations, amounted to 12 sec., costing exactly \$1. Programming time, including typing at a terminal, amounted to about four hours.

This method has been applied to the cases $n = 23$ and $n = 21$.
In each case, an x could not be found such that $S_n(x) < n/2$.

Bibliography

- [1] Diananda, P.H., On a Cyclic Sum, Proc. Glasgow Math. Assoc. 6 (1963), 11-13.
- [2] Djokovic, D. Z., Sur une inégalité, Proc. Glasgow Math. Assoc. 6 (1963), 1-10.
- [3] George, J. Alan, FMIN1 - Local Minimum of a Real Scalar Function of Several Variables Using a Gradient Search Method, Computer Science Department Program Library, Stanford University, 1968.
- [4] Malcolm, M. A. and Katzman, J., RAT - Range Arithmetic Translator, (unpublished), 1969.
- [5] Moore, R. E., Interval Analysis, Prentice-Hall: Englewood Cliffs, N. J., 1966.
- [6] Mordell, L. J., On the Inequality $\sum_{r=1}^n x_r / (x_{r+1} + x_{r+2}) \geq \frac{n}{2}$ and Some Others, Abh. Math. Sem., Univ. of Hamburg 22 (1958), 221-240.
- [7] _____, Note on the Inequality $\sum_{r=1}^n x_r / (x_{r+1} + x_{r+2}) \geq \frac{n}{2}$, J. London Math. Soc. 37 (1962), 176-178.
- [8] Nowosad, Pedro, Isoperimetric Eigenvalue Problems in Algebras, Comm. Pure Appl. Math., 21 (1968), 401-465.
- [9] Rankin, R. A., An Inequality, Math. Gaz. 42 (1958), 39-40.
- [10] Shapiro, H. S., Problem 4603, Amer. Math. Monthly 61 (1954), 571.
- [11] Zulauf, Von A., Note on a Conjecture of L. J. Mordell, Abh. Math. Sem., Univ. of Hamburg 22 (1968), 240-241.